ORIGINAL ARTICLE



Less is more: lemurs (*Eulemur* spp.) may benefit from loss of trichromatic vision

Rachel L. Jacobs¹ · Carrie C. Veilleux² · Edward E. Louis Jr³ · James P. Herrera⁴ · Chihiro Hiramatsu^{5,6} · David C. Frankel¹ · Mitchell T. Irwin⁷ · Amanda D. Melin^{2,8} · Brenda J. Bradley¹

Received: 7 August 2018 / Revised: 12 December 2018 / Accepted: 17 December 2018 / Published online: 25 January 2019 © Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Vertebrate color vision is an ideal system for studying the gains and losses of genetic variation across lineages and impacts on behavior. Among placental mammals, trichromatic vision is unique to primates and is argued to be adaptive for foraging on reddish food. However, trichromacy is variably present in lemurs, including species within the cathemeral genus *Eulemur*, due to inter- and intra-specific variation in X-linked opsin genes. Although this variation could result from genetic drift, it could also reflect ecological adaptation. To understand ecological contributions to color vision variation, we examined cone opsin genes of 11 *Eulemur* species. We found that only *E. flavifrons* and *E. macaco* have polymorphic trichromacy. Most dichromatic species have an "M" (green-shifted) opsin; uniquely, one species (*E. rubriventer*) has dichromatic (M opsin) or polymorphic *Eulemur* ancestor. To address potential ecological explanations for opsin variation, we studied the dietary behavior of wild *E. rubriventer* and collected reflectance spectra from plant species consumed. Visual models suggest that trichromacy should provide an advantage for detecting reddish foods; however, luminance contrasts were greatest for dichromats with the L opsin. As *E. rubriventer* are often active in low-light rainforest conditions, luminance cues may be relatively important, which could favor the L opsin, while also leading to relaxed selection on, or selection against, trichromacy. The presence of different opsin alleles across *Eulemur* species could represent adaptations related to diet, activity pattern, or habitat.

Significance statement

Loss of genetic variation, often thought to be maladaptive, can occur through natural selection. Among primates, some species have trichromatic color vision, the ability to distinguish reddish and greenish hues; others are red-green colorblind (dichromatic).

Amanda D. Melin and Brenda J. Bradley joint senior authors

Electronic supplementary material The online version of this article (https://doi.org/10.1007/s00265-018-2629-9) contains supplementary material, which is available to authorized users.

Rachel L. Jacobs rlynjacobs@gmail.com

- ¹ Center for the Advanced Study of Human Paleobiology, Department of Anthropology, The George Washington University, Washington, DC 20052, USA
- ² Department of Anthropology and Archaeology, University of Calgary, Calgary, AB, Canada
- ³ Conservation Genetics Department, Omaha's Henry Doorly Zoo and Aquarium, Omaha, NE 68107, USA

- ⁴ Department of Evolutionary Anthropology, Duke University, Durham, NC 27708, USA
- ⁵ Department of Human Science, Faculty of Design, Kyushu University, 4-9-1 Shiobaru, Minamiku, Fukuoka 815-8540, Japan
- ⁶ Physiological Anthropology Research Center, Kyushu University, 4-9-1 Shiobaru, Minamiku, Fukuoka 815-8540, Japan
- ⁷ Department of Anthropology, Northern Illinois University, DeKalb, IL 60115, USA
- ⁸ Department of Medical Genetics and Alberta Children's Hospital Research Institute, University of Calgary, Calgary, AB, Canada

We examined adaptive explanations for color vision differences by studying cone opsin genes and behavior in wild lemurs (*Eulemur*)—a genus that is active both day and night. We found that color vision is variable in *Eulemur* species, and full trichromatic vision was likely lost in at least one lineage. Foraging ecology of dichromatic *Eulemur rubriventer* indicates that trichromatic vision should be advantageous for foraging on reddish foods, but brightness cues are more salient to this species' vision. We suggest brightness may be more important than color to this species, particularly at night, and loss of trichromacy could be adaptive in some lemurs.

Keywords Adaptation · Diversity · Luminance · Opsin · Polymorphic trichromacy · Sensory ecology

Introduction

Loss of genetic variation in wild populations is often assumed to be a maladaptive result of genetic drift or inbreeding (Frankham 2005). Indeed, conservation biologists frequently view such losses as an indicator of declining population size and viability (Frankham 1996; Lacy 1997; Reed and Frankham 2003). However, genetic variation can be influenced by multiple evolutionary mechanisms, sometimes pushing allele frequencies in the same direction. For example, adaptive mechanisms like directional selection can also drive the loss of allelic variation (Futuyma 1998), and in a given context "less is more" for some loci. That is, greater allelic variation is not necessarily advantageous and, in some cases, might even be disadvantageous (Futuyma 1998), favoring fewer alleles at a given locus.

The color vision system of vertebrates provides a notable case study in the repeated gain and loss of genetic variation across lineages (e.g., Hunt et al. 1998; Rennison et al. 2012; Borges et al. 2015). Of particular interest, many primates, including nearly all New World monkeys (platyrrhines) and some day-active lemurs, show intra-specific variation in color vision; they have multiple alleles at the single X-linked opsin gene, resulting in "polymorphic trichromacy" (Tan and Li 1999; reviews in Surridge et al. 2003; Veilleux and Bolnick 2009; Kawamura et al. 2012; Jacobs et al. 2017). This pattern differs from other trichromatic primates (e.g., Old World monkeys, apes, and humans), which have two different X-linked opsin genes, and as such, virtually, all individuals have trichromatic color vision (Dulai et al. 1999; Jacobs and Deegan 1999; Nathans 1999; Fig. S1). In species with polymorphic trichromacy, heterozygous females have trichromatic color vision, while homozygous females and all males are redgreen colorblind (i.e., dichromatic; Jacobs and Neitz 1987; Williams et al. 1992; Jacobs et al. 1993; Fig. S1).

Polymorphic color vision in primate populations is often cited as an example of adaptive molecular evolution favoring and maintaining allelic diversity (Surridge et al. 2003; Kawamura and Melin 2017). This system, with its wellestablished genotype-phenotype link (e.g., Bradley and Lawler 2011), provides unique opportunities to address evolutionary questions about how variation is maintained and lost in populations. Accordingly, there has been a large amount of research, particularly in New World monkeys, aimed at identifying the selective mechanisms resulting in balancing selection (e.g., heterozygous advantage, niche differentiation, group benefit of association; Melin et al. 2007, 2008, 2017a; Smith et al. 2012; Veilleux et al. 2016). Although the mechanisms maintaining variation might differ across species and populations, it is reasonable to hypothesize that trichromatic color vision should provide an advantage, or else variation would likely be lost due to allelic drift (Futuyma 1998), and studies of allelic diversity support this argument (Hiwatashi et al. 2010). Until recently, trichromatic advantage remained largely in the realm of theory (e.g., Osorio et al. 2004) with limited evidence for advantages in wild populations. Recent research on New World monkeys, however, has found that trichromatic color vision provides foraging advantages through higher intake rates for trichromats when feeding on vellowish-reddish fruit compared to dichromats (Melin et al. 2017a; but see e.g., Vogel et al. 2007), and more frequent visits to flower patches by trichromatic individuals (Hogan et al. 2018). There is also some evidence that trichromacy provides foraging advantages in polymorphic lemur species (Veilleux et al. 2016). There may be additional advantages to trichromatic individuals related to detecting young leaves (Dominy and Lucas 2001; Melin et al. 2017b), predators (Pessoa et al. 2014), and social signals of conspecifics (Hiramatsu et al. 2017). Taken together, these results support the hypothesis that trichromatic color vision is adaptive and should be favored by natural selection.

What is puzzling then is the potential loss of polymorphic trichromatic color vision in some primates, notably some *Eulemur* species (Jacobs and Bradley 2016). Members of this lineage are cathemeral (active day and night) and eat a large amount of fruit. Although other closely related lemurs with similar biology have polymorphic trichromacy (e.g., genus *Varecia*; Tan and Li 1999; Jacobs and Deegan 2003), it has so far been found in a single species of *Eulemur, E. flavifrons* (Veilleux and Bolnick 2009); other congeners (*E. fulvus, E. collaris, E. mongoz, E. rubriventer*) appear to be dichromatic (Tan and Li 1999; Leonhardt et al. 2009; Jacobs and Bradley 2016; Valenta et al. 2016). Interestingly, different X-linked opsin alleles appear to be fixed in different dichromatic

species. While the polymorphic E. flavifrons has two X-linked opsin alleles (one "M" (green-shifted) opsin with peak spectral sensitivity $[\lambda_{max}] \sim 543$ nm and one "L" (red-shifted) opsin with $\lambda_{\text{max}} \sim 558$ nm; Veilleux and Bolnick 2009; Fig. S1), among the dichromatic congeners, three species (E. fulvus, E. collaris, E. mongoz) exhibit the single M opsin, while one (E. rubriventer) has the L opsin (Tan and Li 1999; Leonhardt et al. 2009; Jacobs and Bradley 2016; Valenta et al. 2016). This distribution of M and L opsins among Eulemur species suggests that polymorphic trichromacy may have been the ancestral Eulemur condition and was subsequently lost in some species or populations (Jacobs and Bradley 2016), a scenario similarly proposed for another dichromatic primate clade, the nocturnal tarsiers (Melin et al. 2013). Importantly, even given a dichromatic Eulemur ancestry (either M or L opsin), polymorphic trichromacy would have been lost in one or more species, as a polymorphic condition, even if brief, must occur to transition between a M and L opsin.

Given that trichromacy appears to be adaptive for foraging on fruit (e.g., Melin et al. 2017a), loss of this variation in *Eulemur* seems surprising (Jacobs and Bradley 2016), because those species are day-active and largely frugivorous (Mittermeier et al. 2010). Allelic loss could be a maladaptive or neutral result of drift (e.g., genetic bottlenecks), but genetic evidence for this mechanism is equivocal (Jacobs and Bradley 2016). We therefore examined whether loss of trichromacy in Eulemur could be adaptive. For example, recent research suggests that fruit colors in a Malagasy forest (Ranomafana National Park) may be less visually conspicuous in the redgreen color channel compared to fruit colors in other forests in Africa (Kibale National Park, Uganda; Nevo et al. 2018). Moreover, although potentially variable across sites (Bollen et al. 2005), those fruits consumed by some lemurs (including some Eulemur species) in Madagascar may be primarily "dull" in coloration (i.e., green and brown; Dew and Wright 1998; Birkinshaw 2001; Bollen et al. 2005). If trichromacy does not provide an advantage in detecting food items consumed by some lemur species/populations, then we might expect selection favoring trichromatic color vision to be relaxed, which could lead to loss of polymorphic trichromacy. Alternatively (or additionally), there could be selection favoring particular opsin variants in dichromatic taxa. Indeed, using color modeling techniques, it has been suggested that L-based dichromacy is superior to M-based dichromacy for detecting chromatic cues of food items consumed by nocturnal woolly lemurs, genus Avahi, in Ranomafana National Park (Veilleux et al. 2014).

Eulemur, therefore, represents an ideal lineage to explore evolutionary mechanisms that could result in loss of variation. The aims of this study were threefold. First, we examined genetic variation across the genus *Eulemur* (11 species and 22 populations) to better characterize the distribution of M and L opsins across populations and species. Second, we explored patterns of potential allele loss in the genus Eulemur by estimating the ancestral color vision state. Finally, we assessed potential ecological pressures related to the visual ecology of dichromatic lemurs, using red-bellied lemurs in Ranomafana National Park as a case study. This population of E. rubriventer is monomorphic for the L opsin (Jacobs and Bradley 2016), and therefore seems to differ from other Eulemur species that are either polymorphic or monomorphic for the M opsin. If trichromacy was lost in this population due to relaxed selection pressures, we predict that the food items consumed will not be more salient to a trichromat compared to a dichromat (Sumner and Mollon 2000; Regan et al. 2001; Hiramatsu et al. 2008). If fixation of the L opsin might represent selection favoring the L opsin over the M opsin during foraging, we predict that the color of food items consumed by E. rubriventer will be more salient to dichromats with the L opsin compared to dichromats with the M opsin. To test these predictions, we measured reflectance spectra of E. rubriventer food items consumed during behavioral observations and then modeled food conspicuity to different color vision systems.

Methods

Study species

Eulemur species are medium-sized frugivorous lemurs that occur throughout Madagascar's varied forest habitats, including the western dry deciduous forests and the eastern rainforests, but they do not occur in the southern spiny forests (Mittermeier et al. 2010). Eulemur species are cathemeral, and species vary in the degree of diurnal and nocturnal activity (Donati et al. 2016). In this study, we examined 11 of the 12 recognized Eulemur species in Madagascar (Fig. 1). We generated opsin sequence data for: E. coronatus, E. flavifrons, E. macaco, E. rubriventer, and members of the so-called brown lemur complex (Markolf et al. 2013), including E. albifrons, E. cinereiceps, E. collaris, E. fulvus, E. rufifrons, and E. sanfordi. Opsin data were already available for captive E. mongoz (Tan and Li 1999). Thus, we examined the entire genus less one species, E. rufus (considered to be part of the "brown lemur complex").

X-linked opsin variation in Eulemur

Sample collection and DNA extraction

Blood/tissue or fecal samples were obtained from 142 individuals across 22 wild populations in Madagascar as part of separate research projects (Table 1; Fig. 1). Blood/tissue samples comprised the majority (N = 138 individuals from 21 populations), and sample collection methods have been

Fig. 1 Ten species of *Eulemur* were sampled across 22 sites in Madagascar. *Identifies Ranomafana National Park where foraging data were collected on red-bellied lemurs (*E. rubriventer*). The "X" under Andringitra indicates a hybrid population (*E. cinereiceps* x *E. rufifrons*). Illustrations copyright 2015 Stephen D. Nash/ IUCN SSC Primate Specialist Group. Used with permission. Map from Du Puy and Moat (1996)



previously described in detail (Brenneman et al. 2012). For those samples, we extracted genomic DNA from blood/tissue using standard phenol-chloroform-isoamyl extraction protocols (Sambrook et al. 1989). Fecal samples were collected from four individuals at the Tsinjoarivo site. Sample collection/preservation methods followed the two-step protocol (ethanol-silica gel beads; Nsubuga et al. 2004), and DNA extraction followed Jacobs and Bradley (2016). Negative controls were included in extractions and all downstream procedures. All genetic analyses were conducted at the George Washington University Primate Genomics Lab and the Conservation Genetics Laboratory at Omaha's Henry Doorly Zoo and Aquarium.

Opsin genotyping

In primates, spectral differences among medium-to-long wavelength sensitive opsins result primarily from amino acid changes from single-nucleotide polymorphisms (SNP) at three sites in the X-linked opsin gene (180 in exon 3; 277 and 285 in exon 5) (Neitz et al. 1991; Hiramatsu et al. 2004). All three sites vary in some lemurs (Jacobs et al. 2017), but among *Eulemur*, only site 285 is known to vary, resulting in two opsin alleles: "M" ($\lambda_{max} \sim 543$ nm with a three-site combination of alanine, tyrosine, alanine—AYA for 180, 277, and 285, respectively) and "L" ($\lambda_{max} \sim 558$ nm with a three-site combination of alanine, tyrosine, threonine— AYT) (Tan and Li 1999; Veilleux and Bolnick 2009). We targeted exons 3 and 5 to capture the three key functional sites.

We amplified exons 3 and 5 using quantitative PCR (Mic qPCR Cycler, Bio Molecular Systems; Rotor-Gene Q, Qiagen), which was immediately followed by high-resolution melt analysis (HRMA; Jacobs et al. 2016). We assigned genotypes for each exon based on the shape and temperature of the melt curve compared to positive controls of known genotypes (identified by Sanger sequencing). Each

 Table 1
 Samples of Eulemur
 individuals for which X-linked opsin genotypes were obtained

Species	Site	$N_{\rm males}, N_{\rm females}$	$N_{\rm X~chromosomes}$	
Eulemur albifrons	Anjanaharibe-Sud	2,3	8	
	Betampona	3,2	7	
	Marojejy	3,2	7	
	Masoala	1,0	1	
		9,7	23	
Eulemur cinereiceps	Andringitra (hybrids)	2,3	8	
	Manombo	3,2	7	
	Vevembe	3,2	7	
		8,7	22	
Eulemur collaris	Andohahela	2,3	8	
	Kalambatritra	3,2	7	
	Midongy du Sud	2,3	8	
		7,8	23	
Eulemur coronatus	Analamerana	3,2	7	
	Andrafiamena	3,0	3	
	Daraina	2,0	2	
	Montagne d'Ambre	3,2	7	
		11,4	19	
Eulemur flavifrons	Sahamalaza	7,3	13	
		7,3	13	
Eulemur fulvus	Andasibe/Mantadia	4,2	8	
	Mariarano	1,4	9	
	Tsinjoarivo	3,0	3	
	Zahamena	3,2	7	
		8,8	24	
Eulemur macaco	Lokobe	1,2	5	
		1,2	5	
Eulemur rubriventer	Andringitra	2,3	8	
	Andasibe	2,3	8	
	Anjanaharibe-Sud	1,1	3	
	Fandriana	3,2	7	
	Marojejy	2,3	8	
	Tsinjoarivo	0,1	2	
	Zahamena	1,0	1	
		11,12	35	
Eulemur rufifrons	Andringitra (hybrids)	3,2	7	
	Isalo	2,3	8	
		5,5	15	
Eulemur sanfordi	Analamerana	1,4	9	
	Andrafiamena	2,0	2	
	Daraina	2,1	4	

sample was replicated in two independent reactions per exon (four reactions per exon for fecal samples). For a subset of individuals of each species (N = 52 for exon 3; N = 79 for exon 5) representing the full range of observed melt temperature

variation (Jacobs et al. 2016), HRMA genotype calls were further confirmed via Sanger sequencing electropherograms of qPCR amplicons. In all cases, sequences matched HRMAbased calls.

3,2

8,7

78,64

Montagne d'Ambre

Total

7

22

206

Ancestral color vision estimations and allele loss

To estimate the evolutionary history of opsin variation in the genus Eulemur, we coded the color vision for 52 species of lemuriforms, lorisiforms, and tarsiiforms as follows: 1-a single M opsin ($\lambda_{max} \leq 543$ nm), 2—polymorphic (two or more alleles), and 3—a single L opsin ($\lambda_{max} \ge 558$ nm) (see Table S1 for data and references). We used the recent phylogeny from Herrera and Dávalos (2016) and added the following missing taxa, based on literature estimating their divergence times: Avahi mooreorum and Tarsius bancanus (Lei et al. 2008; Springer et al. 2012). We mapped the evolution of the three-state character with an ordered transition matrix, such that changing from a single M opsin to a single L opsin or vice versa first required a transition to the polymorphic state. This ordering reflects the dynamics of the visual system. Allelic variation must be present for changes in allele frequencies to occur. Accordingly, changing from a monomorphic M or L opsin genotype (i.e., one opsin allele fixed) to a different monomorphic M or L opsin genotype (i.e., different opsin allele fixed) would necessitate that at some point in the evolutionary history of the population, both opsin alleles were present (e.g., an opsin allele was introduced through mutation or migration), resulting in polymorphic trichromacy. We simulated trait evolution using stochastic character mapping (Huelsenbeck et al. 2003) implemented in the package *phytools* (Revell 2012) for the R statistical environment (R Core Team 2014). Briefly, stochastic character mapping traces the evolution of a discrete trait on a tree using a continuous-time Markov model. This method first calculates the conditional likelihood of character states at each node, then simulates the evolution of the trait along the branches of the tree in proportion to the probability of the trait changes. This approach has the advantage of being a stochastic model of trait evolution that allows more than one state change to occur along branches. The probabilities of state changes on branches are proportional to time; more changes are likely to occur on long branches compared to short branches. The states at ancestral nodes were summarized by averaging, weighted by their probabilities. We ran 10,000 simulations with the ordered transition matrix, estimating the ancestral state at the root under a flat prior probability of 1/3 for each state (make.simmap function in phytools). The analyses were run in two ways: (1) using symmetrical transition rates and (2) allowing transition rates between states to differ.

Foraging ecology of Eulemur rubriventer

We collected foraging data on the population of E. rubriventer

in Ranomafana National Park (RNP), which is an area of

Study site

41,000 ha of montane rainforest in southeastern Madagascar (E47° 18′–47° 37′, S21° 02′–21° 25′; Wright 1992). This study population was previously identified to be monomorphic for the X-linked opsin gene, exhibiting only an L opsin (AYT, $\lambda_{max} \sim 558$ nm; Jacobs and Bradley 2016). Individuals were identifiable for behavioral data collection based on variation in their pelage coloration/patterns (Jacobs and Bradley 2016).

E. rubriventer food items

From the end of September 2012 through mid-May 2013, we collected foraging data on 3 groups of red-bellied lemurs from each of three localities within RNP (N=9)groups; Table 2). It was not possible to record data blind because our study involved focal animals in the field, but behavioral data were collected from focal animals prior to completion of opsin genotyping. We followed groups from the time of group location through dusk until sunset when light levels precluded visual observation of animals. We rotated sites monthly (every 10 days) and attempted to follow each group 3 days/month (Table 2). The goal of the behavioral follows was to characterize the diet of this population (i.e., identify plants, plant parts, and plant part colors that are consumed) based on the foraging behavior of 36 individuals, and thus pseudoreplication was not an issue.

During group follows, we recorded all occurrences (i.e., "bouts") of foraging. We defined bouts as when at least one individual in the group entered a new tree to feed or forage or when foraging resumed in a tree after all individuals had stopped feeding for at least 10 min. During foraging bouts, we recorded the species of the food item consumed (using the local vernacular species names), along with the plant part consumed and the color of the plant part consumed when possible. We defined plant parts as ripe and unripe fruit when visual color changes of the fruit allowed identification (i.e., the use of ripeness category in this study does not refer to quantified mechanical properties of food items). For some fruit, multiple color changes occur during ripening (e.g., green to yellow to red), in which case we assigned a category of "mid-ripe" post hoc based on the color of fruit consumed. For plant species in which the fruit does not exhibit conspicuous color changes during ripening (or if this was unknown), we defined items broadly as "fruit." Additional plant parts included flower buds, flowers, leaf petioles, young leaves, mature leaves, galls, and mushrooms.

Reflectance data

We collected food items consumed by *E. rubriventer* from known feeding trees within 10 days when a study group had been observed feeding. Occasionally ($\sim 15\%$

 Table 2
 Red-bellied lemur study groups, compositions, and number of days followed during behavioral data collection. (AM, adult males; AF, adult females; IM, immature males; IF, immature females)

Group	Locality	$N_{\rm AM}$	$N_{\rm AF}$	$N_{\rm IM}$	$N_{\rm IF}$	N_{Infants}	$N_{\mathrm{Total individuals}}$	$N_{\rm Days}$
TK3	Talatakely	1 (1) ^a	1	1	1	0	4-2	18
TK4	Talatakely	1	1	0	2	0	4	14
TK5	Talatakely	1	1	0	1	0	3	20
VT3	Vatoharanana	1	1	0	0	1	3	16
VT5	Vatoharanana	1	1	1	1	0	4	12
VT7	Vatoharanana	1	1	1	1	0	4	14
VL1	Valohoaka	1	1	1	2	0	5	15
VL5	Valohoaka	1	1	0	1	0	3	18
VL9	Valohoaka	1	1	1	2	0	5	22
		10	9	5	11	1	36	149

^a The group composition of TK3 changed during the course of the study, beginning with four individuals and ending with two individuals (an adult male and adult female). The change in group composition included replacement of the original adult male with a new male

of instances), when known feeding trees were devoid of fruit on collection days, we obtained food items from other trees of the same species within the site. We collected food items directly from trees using an extendable tree pruner when possible. For many trees, however, the height of the tree precluded direct collection, and, in such cases, we collected "fresh" samples (i.e., excluding any overripe or decaying fruit) from the ground (Dominy 2004). Once collected, we placed samples into a cooler with ice packs and returned to the research station and measured the spectral reflectance of food items within 14 h of sample collection (Dominy and Lucas 2004).

We measured reflectance spectra of samples using a USB2000+UV-VIS Miniature Fiber Optic Spectrometer (Ocean Optics, Dunedin, FL) under standard lighting conditions (PX-2 Pulsed Xenon Light Source; Ocean Optics). We recorded measurements relative to a diffuse reflectance standard (WS-1; Ocean Optics) using a reflection probe maintained at a fixed angle (45°) and distance (5 mm) from each sample using a probe holder (RPH-1; Ocean Optics). We frequently recalibrated the spectrometer during data collection to minimize drift. Depending on the size of the sample, we collected multiple (1-5) measurements. In addition to food samples, we collected mature leaves from plant species when possible and recorded one to three measurements for the upper and lower part of each leaf. Food items and leaves were represented by 1-10 (mean = 4) individual samples, and we calculated mean reflectance for each item for data analysis.

Visual modeling analyses

To address the question of whether different color vision phenotypes could provide a foraging advantage, we analyzed the chromaticities and chromatic and achromatic contrasts of food items and calculated visual conspicuousness in units of "Just Noticeable Difference" (JNDs) based on the visual system of *Eulemur*.

Chromaticity analyses We calculated chromaticity using the quantum catch of cone photoreceptors for a trichromatic *Eulemur*: S = 413 nm, M = 543 nm, L = 558 nm (Veilleux and Bolnick 2009; Carvalho et al. 2012). Our calculations followed Hiramatsu et al. (2008) and Valenta et al. (2013, 2016), in which the quantum catch (*Q*) of each cone photoreceptor *i* (i.e., S, M, and L) across 400–700 nm, which represents the visual spectrum of primates, was based on the following formula:

$$Q_i = \int_{400}^{700} R(\lambda) I(\lambda) S_i(\lambda) d(\lambda)$$

In this formula, λ refers to wavelength, R is the reflectance spectrum of the item, I is the spectrum of the illumination, and S is the spectral sensitivity of the cone photoreceptor. We did not include effects of macular pigment on the pre-receptoral filter as this feature is lacking in lemurs. Rather, we calculated functions using methods for lemurs following Valenta et al. (2013, 2016) and included only the effects of the lens.

We used three illumination spectra in our analyses: "day," "dusk," and "moonlit night" (Fig. 2), as we observed red-bellied lemurs to be active and feeding during daylight and dusk (low sun angles 10° to below the horizon; Endler 1993) conditions. Additionally, we include "moonlit night" (unobstructed moonlight) spectra because, although we did not continue group follows throughout the night, *E. rubriventer* has been noted to have extensive nocturnal activity, with two peaks around



Fig. 2 Illumination spectra: "day" (1040 h), "dusk" (1720 h), "moonlit night" (1850 h). The former two were measured in Ranomafana National Park, and the latter was measured in Sabah, Malaysia. Data were

1800-2200 h and 0100-0500 h (Overdorff 1996; Donati et al. 2016). Although some nocturnal conditions preclude the use of color vision, previous research indicates that light conditions at dusk and moonlit conditions at night are likely bright enough to enable color vision in primates (Roth et al. 2008; Melin et al. 2012). The illumination spectrum representing "day" was collected in RNP on April 29, 2013 under light shade (forest canopy) and overcast conditions at 1040 h. The illumination spectrum representing "dusk" was collected near sunset (1720 h) on the same day under open canopy and overcast conditions. These two illumination spectra were measured with down-welling light (probe directed upward) through a cosine corrector (CC-3-DA; Ocean Optics) directly attached to the USB2000+UV-VIS Miniature Fiber Optic Spectrometer. Finally, we used an unobstructed moonlight illumination spectrum taken in the tropical forests of Sabah, Malaysia (July 12, 2011, 1850 h; Melin et al. 2012) as representing a moonlit night. This spectrum was taken with a multichannel spectrometer with a highly sensitive photomultiplier detector and an integrating sphere to ensure a cosine angular response (OL-770VIS, Gooch & Housego, Orlando, FL). Spectra taken in Sabah agree well with irradiance spectra from other latitudes (Endler 1993; Johnsen et al. 2006), indicating that a similar pattern can be expected in Madagascar (cf. Pariente 1980). Irradiance spectra were converted from units of absolute irradiance $(\mu W/m^2/nm)$ to units of

converted to units of photon flux $(\mu mol/m^2/s)$ for visual modeling analyses

photon flux (μ mol/m²/s) prior to calculating chromaticities and JNDs following Maia et al. (2013).

We calculated red-green chromaticity as the ratio of the quantum catch for L cones to L and M cones (i.e., L/(L+M); trichromats only). We calculated blue-yellow chromaticity as the ratio of quantum catch for S cones to L and/or M cones (i.e., S/(L+M) for trichromats; S/M or S/L for dichromats). Finally, we calculated luminance by dividing the quantum catch of L and/or M cones ((L+M) for trichromats; M or L alone for dichromats) by a hypothetical white surface that reflects 100% of the given illumination.

To determine if there is a potential foraging advantage for trichromatic *Eulemur*, we compared chromaticities and luminance of food items to those of mature leaves using non-parametric Mann-Whitney *U* tests. All statistical analyses here and in subsequent sections were performed in R Version 3.4.3 (R Core Team 2017) and were two-tailed with significance set at p < 0.05, followed with Bonferroni correction as appropriate.

Chromatic and luminance contrasts We calculated chromatic and luminance contrasts between each food item consumed and its respective mature leaf background (for upper and lower leaf backgrounds) for dichromatic *Eulemur*. In cases where data for mature leaves of the same species were unavailable, we used mean leaf background (using all mature leaves in the data set) in calculations.

Blue-yellow chromatic contrast was calculated as $|\ln \left(Q_{i_{L,M}}^{f}\right) - \ln \left(Q_{i_{L,M}}^{b}\right)| - |\ln \left(Q_{S}^{f}\right) - \ln \left(Q_{S}^{b}\right)|$. Luminance

contrast was calculated as $|\ln (Q_{i_{L,M}}^f) - \ln (Q_{i_{L,M}}^b)|$. Q is the quantum catch of L or M cones $(i_{L, M})$ for each dichromatic phenotype and S cones (S) for each food item (f) and mature leaf background (b). All calculations followed Hiramatsu et al. (2008) and were performed in Matlab.

To determine if relative chromatic or luminance contrast is greater for dichromats with the L opsin compared to dichromats with the M opsin, we used Wilcoxon signed-rank tests on food items against upper and lower leaf backgrounds for each illumination condition.

JND analyses JND calculations follow established methods (Osorio et al. 2004; Hiramatsu et al. 2008; Matsumoto et al. 2014; Valenta et al. 2016). We show the formula for trichromats below.

JND =
$$\sqrt{\frac{e_S^2 (\Delta f_L - \Delta f_M)^2 + e_M^2 (\Delta f_L - \Delta f_S)^2 + e_L^2 (\Delta f_M - \Delta f_S)^2}{(e_L e_M)^2 + (e_L e_S)^2 + (e_M e_S)^2}}$$

where Δf_i is the difference in the quantum catch of receptor *i* between a food item and its upper or lower leaf background. The noise value (e_i) is set for each receptor type (i = S, M, L) by incorporating both the effect of quantum catch amount and cone proportion in the retina:

$$e_i = \sqrt{\frac{1}{f_i q_i} + \frac{{w_i}^2}{p_i}}$$

where q_i is the estimate of quantum flux (in terms of the number of photons) per cone cell per second in receptor *i*, w_i is the Weber fraction of receptor *i*, and p_i is the relative proportion of receptor *i* to the most abundant cone in the retina (Higham et al. 2010). The relative cone proportions follow Matsumoto et al. (2014) and Valenta et al. (2016). We adjusted maximum photon values for each light environment to represent the decreasing quantity of photons reaching the retina with lowering ambient light: $q_{L, M}$ = 10^4 ("day"), $q_{L, M} = 10^3$ ("dusk"), $q_{L, M} = 10^2$ ("moonlit night"). We set the value of quantum flux for S cones as $q_s = q_I/10$ to take account of their low sensitivity. We used Weber fraction values (w_i) of 0.08 for the S cones and 0.02 for both the M and the L cones. These values are close to psychophysical thresholds for humans (Wyszecki and Stiles 1982; Osorio et al. 2004). In the absence of species-specific data on relative cone proportions in the retinas, we set p_L , $p_M = 1$, $p_s = 0.1$ (Martin and Grunert 1999; Hiramatsu et al. 2008). JND values were calculated for each Eulemur color vision phenotype, and we assessed differences using Wilcoxon signed-rank tests.

Data availability Opsin sequence data generated during this study are available as electronic supplementary files.

Reflectance spectra analyzed during the study are available from the corresponding author on reasonable request.

Results

X-linked opsin variation in Eulemur

Opsin genotyping

Results of our opsin analyses identified variation in the presence of opsin alleles across Eulemur species, but not across populations within species (Fig. 3). Seven species of Eulemur were monomorphic for the M opsin (no L allele): E. albifrons, E. cinereiceps, E. collaris, E. coronatus, E. fulvus, E. rufifrons, and E. sanfordi. All sequence traces had the following amino acid combination for sites 180, 277, and 285, respectively: alanine, tyrosine, alanine (AYA, $\lambda_{max} \sim 543$ nm). All highresolution melt curves shared similar shape and temperature profiles as those samples that were sequenced (N = 44 for exon)3; N = 65 for exon 5), indicating no variation. In contrast, all individuals in all populations of red-bellied lemurs (E. rubriventer) were monomorphic for the L opsin (no M allele). Sequence traces (N = 4 exon 3; N = 8 exon 5) showed the following three-site combination: alanine, tyrosine, threonine (AYT, $\lambda_{max} \sim 558$ nm). All samples exhibited similar shape and temperature profiles, indicating no variation. Both M and L alleles (AYA and AYT), and thus polymorphic trichromacy, were only found in E. flavifrons and E. macaco. These were confirmed by sequence traces (N = 4 for exon 3; N = 6 for exon 5) and high-resolution melt curves. Consensus sequences for each species are available in ESM 2 and ESM 3 (for polymorphic species, the A-G nucleotide polymorphism is coded as R).

Ancestral color vision estimation and allele loss

The results of the stochastic character mapping analyses using two transition-rate patterns (symmetrical rates and different rates) are similar (Fig. S2). Accordingly, we report results for the analysis allowing transition rates to differ. Our results support either the M opsin or polymorphic trichromacy (M and L opsin) as the ancestral state for the genus *Eulemur* (posterior probability of M opsin only = 0.72, posterior probability of polymorphic trichromacy = 0.28, Fig. 3). Loss of polymorphic trichromacy likely occurred in one (*E. rubriventer*) or more *Eulemur* species.

Polymorphic trichromacy was the most likely ancestral state for the family Indriidae and the genus *Propithecus*, with apparent M opsin loss in the nocturnal genus *Avahi* (Fig. S2). Deeper nodes in the tree are estimated with less certainty, either as the M opsin or polymorphic. For example, the ancestral state for all Lemuridae has near-equal posterior probabilities of being either monomorphic with an M opsin (0.48) or

Fig. 3 Opsin gene variation and ancestral state estimations based on stochastic character mapping analyses for the genus Eulemur. Green circles at the tips represent the presence of the M opsin only $(\lambda_{max} \sim 543 \text{ nm})$, the red patterned circle represents the presence of the L opsin only (E. rubriventer), $(\lambda_{\text{max}} \sim 558 \text{ nm})$, and gray circles at the tips represent the presence of both M and L opsins (polymorphic trichromacy). Pie charts at each node represent the posterior probabilities for cone opsin ancestral states (colors defined as for the tips). Phylogeny from Herrera and Dávalos (2016). *Indicates data from published material (Tan and Li 1999). Illustrations copyright 2015 Stephen D. Nash/IUCN SSC Primate Specialist Group. Used with permission



polymorphic (0.47; posterior probability of L opsin only = 0.05). State changes most frequently occurred from the polymorphic state to the M opsin (10.6 transitions, on average across 10,000 simulations) or vice versa (8.93 transitions), and less frequently from the polymorphic state to the L opsin (8.45 transitions) or from the L opsin to the polymorphic state (4.27). Across the tree, there were on average 19 independent losses of an opsin allele. Lineages spent 51% of the time in the M opsin state, 31% of the time in the polymorphic state.

Foraging ecology of Eulemur rubriventer

E. rubriventer food items

We recorded a total of 2924 foraging bouts on plant material during the study period. Table S2 lists all species and plant parts consumed, as well as the percentage of foraging bouts

for each plant taxon. Overall, the nine groups of *E. rubriventer* fed on 115 plant taxa. Fruit foraging accounted for the majority of bouts (1947; 67%), followed by flowers/flower buds (480; 16%), and leaves (399; 14%). For the fruit foraging bouts for which ripeness of food items consumed could be determined (including N = 58 species), unripe fruit accounted for 56% of bouts.

Visual modeling analyses

We included reflectance data for 40 species (72 plant parts; see Fig. S3, for example, spectra) consumed by RNP red-bellied lemurs in our analyses (Table S2). The 40 species represent 75% of observed foraging bouts.

Chromaticity analyses Chromaticity plots for a trichromatic *Eulemur* under "day," "dusk," and "moonlit night" conditions are illustrated in Fig. 4 (see also Fig. S4). Red-green (L/(L+

Fig. 4 Chromaticity and luminance plots under "day," "dusk," and "moonlit night" illuminations for 72 plant parts from 40 plant species consumed by *E. rubriventer* in RNP. Mean value is plotted for each plant part. Left, red-green (L/(L+M)) chromaticity vs. blue-yellow (S/ (L + M)) chromaticity plots; right, luminance (L+M) vs. blue-yellow (S/(L+M)) chromaticity plots; S opsin $\lambda_{max} = 413$ nm, M opsin $\lambda_{max} = 543$ nm, L opsin $\lambda_{max} = 558$ nm



M)) vs. blue-yellow (S/(L+M)) chromaticity plots reveal that most food items have greater red-green chromaticity compared to mature leaves under all illumination conditions. Results of Mann-Whitney *U* tests reveal that red-green chromaticities of food items are significantly greater than mature leaves under all illuminations ("day," upper leaf W=2084, p < 0.001; lower leaf, W=1741, p < 0.001; "dusk," upper leaf, W=1974, p < 0.001; lower leaf, W=1613, p < 0.001; "moonlit night," upper leaf, W=2086, p < 0.001; lower leaf, W=1783, p < 0.001; Fig. 5; See Table S3 for descriptive statistics). Results hold under a Bonferroni corrected significance level (α < 0.008).

Our results suggest that many food items with greater redgreen chromaticities are ripe fruit (Fig. S4). To explore this further, we performed a post hoc Mann-Whitney U test to determine if ripe fruit (N=21) has significantly greater redgreen chromaticity than all other food items (N=51). We found that the red-green chromaticity of ripe fruit is greater than that of other food items under "day" (W = 232, p < 0.001), "dusk" (W = 257, p < 0.001), and "moonlit night" (W = 223, p < 0.001) conditions.

Luminance (L+M) vs. blue-yellow chromaticity plots indicate that food items largely overlap with mature leaves in luminance as well as in blue-yellow chromaticity (Fig. 4, Fig. S4). A similar pattern of overlap in luminance and blueyellow chromaticity for a trichromatic lemur is found in both dichromatic phenotypes (Fig. S5). Despite the large amount of overlap apparent from the chromaticity plots, for a trichromatic *Eulemur*, blue-yellow chromaticity is also significantly greater for food items compared to mature upper leaves ("day," W = 1453, p < 0.01; "dusk," W = 1492, p < 0.001; "moonlit night," W = 1393, p < 0.01). However, results do not hold for "moonlit night" conditions under Bonferroni correction ($\alpha < 0.008$). Compared to lower leaves, blue-yellow chromaticity is significantly greater under some light levels ("day," W = 1367, p < 0.05; "dusk," W = 1405, p < 0.01;



Fig. 5 Box-and-whiskers plots of red-green chromaticities (L/(L+M)) of food items, upper leaves (UL), and lower leaves (LL) under the three illumination conditions modeled in this study ("day," "dusk," and "moonlit night"); M opsin $\lambda_{max} = 543$ nm, L opsin $\lambda_{max} = 558$ nm.

Horizontal lines within each box indicate the median of the distribution. Boxes envelop the interquartile range (50% of values) of the sample distribution, and whiskers encompass 1.5 times the interquartile range

"moonlit night," W = 1301, p = 0.054), but results only hold for "dusk" under Bonferroni correction ($\alpha < 0.008$). Luminance is significantly greater for food items compared to mature upper leaves ("day," W = 1842, p < 0.001; "dusk," W = 1824, p < 0.001; "moonlit night," W = 1902, p < 0.001), but not lower leaves ("day," W = 1239, p = 0.144; "dusk," W = 1214, p = 0.203; "moonlit night," W = 1296, p = 0.059; Bonferroni correction, $\alpha < 0.008$).

Contrast analyses Chromatic and luminance contrasts of food items differ between the two dichromatic phenotypes (Tables 3 and 4). Blue-yellow chromatic contrasts are greater for dichromats with the M opsin than chromatic contrasts based on the L opsin. This pattern holds under "day," "dusk," and "moonlit night" conditions (N = 72; upper leaf and lower leaf (all conditions), p < 0.001; see Table 3 for descriptive statistics), as well as under Bonferroni correction ($\alpha <$ 0.008). Luminance contrasts, on the other hand, are significantly greater for dichromats with the L opsin compared to dichromats with the M opsin under all illuminants when viewed under upper leaf conditions (N = 72; upper leaf (all conditions), p < 0.001; Table 4), but not lower leaf conditions ("day," V = 1168, p = 0.414; "dusk," V = 1336, p = 0.904; "moonlit night," V = 1092; p = 0.214; Bonferroni correction, $\alpha < 0.008).$

JND analyses Results of the JND analyses indicate that a higher proportion of food items exhibit ≥ 1 chromatic JND value for trichromatic *Eulemur* compared to dichromatic *Eulemur* under "day" and "dusk" conditions. The proportions of food items that exhibit ≥ 1 chromatic JND value are similarly low ($\leq 10\%$) for all color vision phenotypes under "moonlit night" conditions (Table 5). The proportions of food

items with ≥ 1 chromatic JND value are similar for both dichromatic phenotypes, but dichromats with the L opsin exhibit the lowest proportions under all illumination conditions. Most food items ($\geq 60\%$) have ≥ 1 luminance JND value for all color vision phenotypes, and proportions are similar across phenotypes under each illumination condition (Table 5).

Trichromats have significantly higher chromatic JND values compared to dichromats with the M opsin ("day," upper leaf, V = 22, p < 0.001; lower leaf, V = 99, p < 0.001; "dusk," upper leaf, V = 8, p < 0.001; lower leaf, V = 0; p < 0.001; "moonlit night," upper leaf, V = 2, p < 0.001; lower leaf, V = 1, p < 0.001) and dichromats with the L opsin ("day," upper leaf, V = 0, p < 0.001; lower leaf, V = 4, p < 0.001; "dusk," upper leaf, V = 0, p < 0.001; lower leaf, V = 8; p < 0.001; "moonlit night," upper leaf, V = 0, p < 0.001; lower leaf, V = 0, p < 0.001) (Fig. S6). Dichromats with the M opsin have significantly higher chromatic JND values compared to dichromats with the L opsin ("day," upper leaf, V = 1963, p < 0.001; lower leaf, V = 2056, p < 0.001; "dusk," upper leaf, V = 1939, p < 0.001; lower leaf, V = 1860; p < 0.01; "moonlit night," upper leaf, V = 1992, p < 0.001; lower leaf, V = 2011, p < 0.001) (Fig. S6). All results hold under Bonferroni correction ($\alpha < 0.0028$).

Luminance JND values (Fig. S7) are significantly greater under upper leaf conditions for dichromats with the L opsin compared to trichromats ("day," V = 2284, p < 0.001; "dusk," V = 2269, p < 0.001, "moonlit night," V = 2521, p < 0.001), and compared to dichromats with the M opsin ("day," V =396, p < 0.001; "dusk," V = 372, p < 0.001; "moonlit night," V = 124, p < 0.001) (Fig. S7). Luminance JND values are significantly greater for trichromats compared to dichromats with the M opsin under upper leaf conditions ("day," V =444, p < 0.001; "dusk," V = 432, p < 0.001; "moonlit night," V = **Table 3** Descriptive statistics of blue-yellow chromatic contrasts offood items (N=72) against their leaf backgrounds (UL, upper leaves;LL, lower leaves) for the two *Eulemur* dichromatic phenotypes. Dataare presented for the three illumination conditions modeled in this study.

Background	Dichromat L opsin							Dichromat M opsin					
	Mean	SD	SE	25th %	Median	75th %	Mean	SD	SE	25th %	Median	75th %	
Day													
UL	0.082	0.072	0.009	0.027	0.061	0.119	0.095	0.086	0.010	0.028	0.067	0.126	
LL	0.071	0.069	0.008	0.023	0.048	0.094	0.084	0.081	0.010	0.029	0.059	0.103	
Dusk													
UL	0.091	0.079	0.009	0.029	0.073	0.129	0.107	0.096	0.011	0.034	0.074	0.143	
LL	0.079	0.074	0.009	0.026	0.057	0.108	0.095	0.090	0.011	0.033	0.068	0.129	
Moonlit night													
UL	0.074	0.067	0.008	0.023	0.052	0.113	0.085	0.077	0.009	0.026	0.064	0.121	
LL	0.065	0.063	0.007	0.022	0.045	0.083	0.075	0.073	0.009	0.024	0.050	0.095	

221, p < 0.001). Significant results hold under Bonferroni correction ($\alpha < 0.0028$).

Luminance JND values are not significantly different for trichromats under lower leaf conditions compared to each dichromatic phenotype (M opsin, "day," V = 1177, p = 0.444; "dusk," V = 1152, p = 0.365; "moonlit night," V = 1025, p = 0.106; L opsin, "day," V = 1530, p = 0.227; "dusk," V = 1561, p = 0.167; "moonlit night," V = 1637, p = 0.070), nor are they significantly different between dichromats ("day," V = 1140, p = 0.330; "dusk," V = 1124; p = 0.288; "moonlit night," V = 1003, p = 0.081) (Fig. S7).

Discussion

Our results indicate that extant *Eulemur* species differ in color vision capacity, and only two species have a cone opsin

polymorphism. Wild populations of E. flavifrons exhibit polymorphic trichromacy (as previously reported for a captive population; Veilleux and Bolnick 2009) as does its sister species, E. macaco (Fig. 3). Most other species of Eulemur appear to be monomorphic for the M opsin. E. rubriventer differs from all other congeners in being monomorphic for the L opsin. Variable distribution of opsin alleles occurs throughout the lemur tree and among closely related lemurids (i.e., Varecia-polymorphic, Hapalemur-L opsin, Lemur-M opsin; Tan and Li 1999), which might suggest a polymorphic ancestry in the genus Eulemur. However, our ancestral state estimations are more equivocal, with a higher probability of ancestral Eulemur being monomorphic for the M opsin based on a recent phylogeny. Together, these results suggest that polymorphic trichromacy was likely lost in at least one Eulemur species: E. rubriventer. Interestingly, our visual modeling analyses suggest that trichromatic color vision

Table 4Descriptive statistics of luminance contrasts of food items (N =72) against their leaf backgrounds (UL, upper leaves; LL, lower leaves)for the two *Eulemur* dichromatic phenotypes. Data are presented for the

three illumination conditions modeled in this study. For reference, the luminance contrast of a "reddish" ripe fruit (guava, Fig. S3) for a trichromatic phenotype under upper leaf and "day" conditions is 0.061

Background	Dichromat L Opsin							Dichromat M Opsin					
	Mean	SD	SE	25th %	Median	75th %	Mean	SD	SE	25th %	Median	75th %	
Day													
UL	0.425	0.262	0.031	0.227	0.429	0.584	0.399	0.252	0.030	0.219	0.384	0.563	
LL	0.247	0.181	0.021	0.093	0.243	0.355	0.252	0.190	0.022	0.080	0.231	0.371	
Dusk													
UL	0.415	0.260	0.031	0.227	0.414	0.579	0.397	0.253	0.030	0.202	0.389	0.551	
LL	0.249	0.188	0.022	0.095	0.234	0.354	0.257	0.199	0.023	0.094	0.231	0.388	
Moonlit night													
UL	0.457	0.263	0.031	0.236	0.478	0.609	0.414	0.258	0.030	0.217	0.399	0.579	
LL	0.247	0.181	0.021	0.116	0.223	0.357	0.248	0.183	0.022	0.087	0.233	0.355	

Table 5 Proportions of food items exhibiting ≥ 1 JND for each colorvision genotype under each illumination condition and for upper andlower leaves (upper/lower). Results are presented for chromatic andluminance JNDs

Illumination	Trichromat	Dichromat M opsin	Dichromat L opsin
Chromatic JND			
Day	82/76	67/65	63/60
Dusk	47/49	38/36	33/35
Moonlit night	10/3	10/3	7/0
Luminance JND)		
Day	94/92	94/90	93/90
Dusk	92/79	90/82	89/83
Moonlit night	78/64	76/63	81/65

would likely provide a foraging advantage to E. rubriventer during daylight and dusk conditions, which would seem to favor maintaining a cone opsin polymorphism. Although chromatic values for food items are comparatively higher for trichromatic Eulemur under all conditions, the proportions of food items that are chromatically conspicuous to *Eulemur* are similarly low for trichromatic and dichromatic individuals under moonlit night, indicating that trichromatic color vision is unlikely advantageous under nocturnal conditions. Moreover, we found that many food items should be conspicuous to dichromats in the absence of red-green color vision. Accordingly, given the cathemeral behavior of E. rubriventer, trichromatic color vision might not be under strong selection in this taxon, and fixation of the L opsin could therefore result from relaxed selection (genetic drift; Jacobs and Bradley 2016).

Our visual modeling analyses revealed intriguing differences between the two dichromatic phenotypes that could result in directional selection favoring the L opsin. Specifically, chromatic contrasts and chromatic JND values for *E. rubriventer* food items are significantly greater for dichromats with the M opsin compared to those with only the L opsin. On the other hand, luminance contrasts and luminance JND values are generally greater for dichromats with the L opsin compared to dichromats with the M opsin. Moreover, luminance JND values are generally greater for dichromats with the L opsin compared to trichromats. Therefore, we hypothesize that luminance is a more important foraging cue to *E. rubriventer*, favoring the adaptive L opsin.

Such an adaptation could be related to the particular diet of *E. rubriventer*, but other factors, such as activity pattern, might also play a role. For example, luminance vision may be particularly important under low light level conditions at night when chromatic conspicuity of food items is greatly reduced and the use of color vision may be precluded. *E. rubriventer* appears to have the lowest proportion of diurnal activity among cathemeral *Eulemur* for which data are

available (Donati et al. 2016), which is consistent with this species relying more heavily on luminance vision. If luminance vision is highly relevant to *E. rubriventer*, then the L opsin, being superior for luminance vision compared to the M opsin, may have been fixed by natural selection.

In line with this hypothesis, E. rubriventer is restricted to rainforest environments, which under moonlit conditions appear to be richer in longer wavelengths compared to Madagascar's dry forest environments (Veilleux and Cummings 2012). The L opsin might therefore maximize photon absorption under these nocturnal rainforest conditions and thus allow for better luminance vision. Madagascar's dry forests, on the other hand, are comparatively richer in shorter and middle wavelengths under moonlight (Veilleux and Cummings 2012). Interestingly, unlike E. rubriventer, E. mongoz, some populations of E. coronatus, and at least some species of the "brown lemur complex" inhabit dry forests (Mittermeier et al. 2010). Although the M opsin appears to be the more likely ancestral Eulemur condition based on our analyses, indicating this trait may have been maintained in these lineages, it could also be adaptive for luminance vision under Madagascar's dry forest environments. Dichromatic color vision and fixation of different opsin alleles might therefore represent adaptations to nocturnal activity in different habitats. Resolving the ancestral color vision states and patterns of allele loss throughout the lemur lineage could help address this hypothesis.

If E. rubriventer color vision is related to a greater reliance on luminance vision, this presents another, non-mutually exclusive, hypothesis for potential loss of polymorphic trichromacy. Specifically, loss could result from selection against trichromacy. Previous research indicates that chromatic information corrupts luminance vision (Osorio et al. 1998). Although this is not a hypothesis we could formally test, as the effect is not accounted for in our visual modeling analyses, it likely explains why dichromatic primates exhibit greater foraging efficiency than trichromats on some camouflaged food items (Melin et al. 2007, 2010; Caine et al. 2010; Smith et al. 2012). For species that rely heavily on luminance vision, dichromacy may therefore have a net advantage leading to disruptive selection (i.e., selection against trichromacy). This raises the question why cathemeral E. flavifrons and E. macaco have polymorphic trichromacy, despite potential costs to luminance vision. As has been suggested by other researchers (Valenta et al. 2016), opsin variation could be related to differences in diet and/or activity patterns. For example, polymorphic trichromacy might be favored in species that have a high proportion of daytime activity (Valenta et al. 2016), or that spend large amounts of time foraging under high light levels (Yamashita et al. 2005). Although current comparative research does not necessarily support the former hypothesis (Donati et al. 2016), a population of polymorphic Propithecus verreauxi appears to feed at higher light levels compared to sympatric dichromatic Lemur catta (Yamashita et al. 2005). Interestingly, the two polymorphic *Eulemur* species are known to inhabit a unique transitional environment between eastern rainforests and western dry forests (Sambirano) (Mittermeier et al. 2010), but whether this environment imposes a unique selective pressure is unknown.

As a final note, it is important to acknowledge that the psychophysical data available for lemur vision are limited. Chromatic and achromatic discrimination thresholds and data on cone ratios are currently lacking for lemurs (Olsson et al. 2018). These parameters can have a significant impact on the models used in our study (Olsson et al. 2018), which were designed for understanding chromatic processing separate from achromatic processing (Osorio and Vorobyev 2018). Accordingly, our results of the modeling analyses should be interpreted with caution and considered hypotheses to be tested as our understanding of lemur visual processing improves.

Moving forward, what is clearly needed, in addition to the psychophysical information above, is to examine the detailed foraging behaviors and visual environments of different *Eulemur* species/populations to test whether opsin variation seen across species is likely adaptive or neutral. Together with behavioral studies, determining whether (1) similar patterns in chromatic versus luminance conspicuity are observed for plant species consumed by dichromatic lemurs with the M opsin, and (2) trichromacy is advantageous to individual *E. flavifrons* and *E. macaco* would be highly instructive for assessing the evolutionary mechanisms underlying differences in color vision capacities.

Although we cannot definitively conclude that loss of opsin variation is adaptive for some lemurs, such as *E. rubriventer*, the results of this study underscore the importance of considering the many drivers of allelic variation and allele loss when assessing heterozygosity in conservation genetics. Loss of variation can potentially be adaptive and thus external efforts to increase diversity (e.g., via outcrossings; Johnson et al. 2010) could yield unintended consequences.

Acknowledgements We thank Benjamin Andriamihaja and MICET, Ministère de l'Environnement, de l'Écologie et des Forêts, Madagascar National Parks, Eileen Larney and the Centre ValBio, and the University of Antananarivo for providing logistical support and research permissions in Madagascar. We thank S. Ambler, C. Angyal, B. Chowdhury, J. Falinomenjanahary, J.P. Lahitsara, A. Minoasy, N. Phelps, T. Randriarimanga, E. Razafimandimby, D. Razafindraibe, J. Razafindramasy, A. Telo, A.V. Tombotiana, Velomaharavo, and J.B. Velontsara for assisting with data collection in Madagascar. We also thank Stephen D. Nash for generously allowing use of his illustrations in this manuscript's Figs. We thank Lauren Anderson, Melissa T. R. Hawkins, Cynthia Frasier, Shannon Engberg, and Carolyn Bailey for assistance with lab analyses. Finally, we thank three anonymous reviewers for their helpful comments.

Funding This research was funded in part by the National Science Foundation (DDIG, BCS 1232535), the Leakey Foundation, the Wenner-Gren Foundation, the RW Primate Fund, Yale University, The George Washington University, and the Interdepartmental Doctoral Program in Anthropological Sciences at Stony Brook University.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All behavioral protocols and animal handling procedures were approved by and adhered to institutional animal care requirements (Stony Brook IACUC# 2011-1895, Omaha's Henry Doorly Zoo and Aquarium IACUC# 97-001, 12-101, and Northern Illinois University IACUC #LA12-0011) and national laws. Data collection, sample collection, and export permits were obtained from Madagascar National Parks, formerly Association Nationale pour la Gestion des Aires Protégées (ANGAP), and the Ministère de l'Environnement, de l'Écologie et des Forêts. Samples were exported/imported under the Convention on International Trade in Endangered Species (CITES) Appendix I permits.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Birkinshaw C (2001) Fruit characteristics of species dispersed by the black lemur (*Eulemur macaco*) in the Lokobe Forest, Madagascar. Biotropica 33:478–486
- Bollen A, Donati G, Fietz J, Schwab D, Ramanamanjato J-P, Randrihasipara L, van Elsacker L, Ganzhorn J (2005) An intersite comparison of fruit characteristics in Madagascar: evidence for selection pressure through abiotic constraints rather than through coevolution. In: Dew JL, Boubli JP (eds) Tropical fruits and frugivores: the search for strong interactors. Springer, The Hague, pp 93–119
- Borges R, Khan I, Johnson WE, Gilbert MTP, Zhang G, Jarvis ED, O'Brien SJ, Antunes A (2015) Gene loss, adaptive evolution and the co-evolution of plumage coloration genes with opsins in birds. BMC Genomics 16:751
- Bradley BJ, Lawler RR (2011) Linking genotypes, phenotypes, and fitness in wild primate populations. Evol Anthropol 20:104–119
- Brenneman RA, Johnson SE, Bailey CA, Ingraldi C, Delmore K, Wyman TM, Andriamaharoa HE, Ralainasolo FB, Ratsimbazafy JH, Louis EE Jr (2012) Population genetics and abundance of the endangered grey-headed lemur *Eulemur cinereiceps* in south-east Madagascar: assessing risks for fragmented and continuous populations. Oryx 46: 298–307
- Caine NG, Osorio D, Mundy NI (2010) A foraging advantage for dichromatic marmosets (*Callithrix geoffroyi*) at low light intensity. Biol Lett 6:36–38
- Carvalho LS, Davies WL, Robinson PR, Hunt DM (2012) Spectral tuning and evolution of primate short-wavelength-sensitive visual pigments. Proc R Soc Lond B 279:387–393
- Dew JL, Wright P (1998) Frugivory and seed dispersal by four species of primates in Madagascar's eastern rain forest. Biotropica 30:425–437
- Dominy NJ (2004) Fruits, fingers, and fermentation: the sensory cues available to foraging primates. Integr Comp Biol 44:295–303
- Dominy NJ, Lucas PW (2001) Ecological importance of trichromatic vision to primates. Nature 410:363–366
- Dominy NJ, Lucas PW (2004) Significance of color, calories, and climate to the visual ecology of catarrhines. Am J Primatol 62:189–207
- Donati G, Campera M, Balestri M, Serra V, Barresi M, Schwitzer C, Curtis DJ, Santini L (2016) Ecological and anthropogenic correlates of activity patterns in *Eulemur*. Int J Primatol 37:29–46
- Du Puy DJ, Moat J (1996) A refined classification of the primary vegetation of Madagascar based on the underlying geology: using GIS to map its distribution and to assess its conservation status. In:

Lourenço WR (ed) Proceedings of the International Symposium on the Biogeography of Madagascar. Editions de l'ORSTOM, Paris, pp 205–218 + 3 maps

- Dulai KS, von Dornum M, Mollon JD, Hunt DM (1999) The evolution of trichromatic color vision by opsin gene duplication in New World and Old World primates. Genome Res 9:629–638
- Endler JA (1993) The color of light in forests and its implications. Ecol Monogr 63:1–27
- Frankham R (1996) Relationship of genetic variation to population size in wildlife. Conserv Biol 10:1500–1508
- Frankham R (2005) Genetics and extinction. Biol Conserv 126:131–140
- Futuyma DJ (1998) Evolutionary biology, 3rd edn. Sinauer Associates Inc, Sunderland
- Herrera JP, Dávalos LM (2016) Phylogeny and divergence times of lemurs inferred with recent and ancient fossils in the tree. Syst Biol 65: 772–791
- Higham JP, Brent LJN, Dubuc C, Accamando AK, Engelhardt A, Gerald MS, Heistermann M, Stevens M (2010) Color signal information content and the eye of the beholder: a case study in the rhesus macaque. Behav Ecol 21:739–746
- Hiramatsu C, Radlwimmer FB, Yokoyama S, Kawamura S (2004) Mutagenesis and reconstitution of middle-to-long-wave-sensitive visual pigments of New World monkeys for testing the tuning effect of residues at sites 229 and 233. Vis Res 44:2225–2231
- Hiramatsu C, Melin AD, Aureli F, Schaffner CM, Vorobyev M, Matsumoto Y, Kawamura S (2008) Importance of achromatic contrast in short-range fruit foraging in primates. PLoS One 3:e3356
- Hiramatsu C, Melin A, Allen W, Dubuc C, Higham J (2017) Experimental evidence that primate trichromacy is well suited for detecting primate social colour signals. Proc R Soc B 284:20162458
- Hiwatashi T, Okabe Y, Tsutsui T, Hiramatsu C, Melin AD, Oota H, Schaffner CM, Aureli F, Fedigan LM, Innan H, Kawamura S (2010) An explicit signature of balancing selection for colour vision variation in New World monkeys. Mol Biol Evol 27:453–464
- Hogan J, Fedigan L, Hiramatsu C, Kawamura S, Melin A (2018) Florivory reveals detection advantage of small, ephemeral resources to trichromatic New World monkeys. Sci Rep 8:10883
- Huelsenbeck JP, Nielsen R, Bollback JP (2003) Stochastic mapping of morphological characters. Syst Biol 52:131–158
- Hunt DM, Dulai KS, Cowing JA, Julliot C, Mollon JD, Bowmaker JK, Li WH, Hewett-Emmett D (1998) Molecular evolution of trichromacy in primates. Vis Res 38:3299–3306
- Jacobs RL, Bradley BJ (2016) Considering the influence of nonadaptive evolution on primate color vision. PLoS One 11:e0149664
- Jacobs GH, Deegan JF (1999) Uniformity of colour vision in Old World monkeys. Proc R Soc Lond B 266:2023–2028
- Jacobs GH, Deegan JF (2003) Photopigment polymorphism in prosimians and the origins of primate trichromacy. In: Mollon JD, Pokorny J, Knoblauch K (eds) Normal and defective colour vision. Oxford University Press, Oxford, pp 14–20
- Jacobs GH, Neitz J (1987) Inheritance of color vision in a New World monkey (*Saimiri sciureus*). Proc Natl Acad Sci U S A 84:2545– 2549
- Jacobs GH, Neitz J, Neitz M (1993) Genetic basis of polymorphism in the color vision of platyrrhine monkeys. Vis Res 33:269–274
- Jacobs RL, Spriggs AN, MacFie TS, Baden AL, Irwin MT, Wright PC, Louis EE, Lawler RR, Mundy NI, Bradley BJ (2016) Primate genotyping via high resolution melt analysis: rapid and reliable identification of color vision status in wild lemurs. Primates 57:541–547
- Jacobs RL, MacFie TS, Spriggs AN et al (2017) Novel opsin gene variation in large-bodied, diurnal lemurs. Biol Lett 13:20170050
- Johnsen S, Kelber A, Warrant E, Sweeney AM, Widder EA, Lee RL, Hernández-Andrés J (2006) Crepuscular and nocturnal illumination and its effects on color perception by the nocturnal hawkmoth *Deilephila elpenor*. J Exp Biol 209:789–800

- Johnson WE, Onorato DP, Roelke ME, Land ED, Cunningham M, Belden RC, McBride R, Jansen D, Lotz M, Shindle D, Howard J, Wildt DE, Penfold LM, Hostetler JA, Oli MK, O'Brien SJ (2010) Genetic restoration of the Florida panther. Science 329:1641–1645
- Kawamura S, Melin AD (2017) Evolution of genes for color vision and the chemical senses in primates. In: Saitou N (ed) Evolution of the human genome I. Springer, Tokyo, pp 181–216
- Kawamura S, Hiramatsu C, Schaffner CM, Melin AD, Aureli F, Fedigan LM (2012) Polymorphic color vision in primates: evolutionary considerations. In: Hirai H, Imai H, Go Y (eds) Post-genome biology of primates. Springer, Tokyo, pp 93–120
- Lacy RC (1997) Importance of genetic variation to the viability of mammalian populations. J Mammal 78:320–335
- Lei R, Engberg SE, Andriantompohavana R, McGuire SM, Mittermeier RA, Zaonarivelo JR, Brenneman RA, Louis EE Jr (2008) Nocturnal lemur diversity at Masoala National Park. Spec Pub MusTexas Tech Univ 53:1–48
- Leonhardt SD, Tung J, Camden JB, Leal M, Drea CM (2009) Seeing red: behavioral evidence of trichromatic color vision in strepsirrhine primates. Behav Ecol 20:1–12
- Maia R, Eliason C, Bitton P, Doucet S, Shawkey M (2013) pavo: an R package for the analysis, visualization and organization of spectral data. Methods Ecol Evol 4:906–913
- Markolf M, Rakotonirina H, Fichtel C, von Grumbkow P, Brameier M, Kappeler PM (2013) True lemurs...true species – species delimitation using multiple data sources in the brown lemur complex. BMC Evol Biol 13:233
- Martin PR, Grunert U (1999) Analysis of short wavelength sensitive ("blue") cone mosaic in the primate retina: a comparison of New World and Old World monkeys. J Comp Neurol 406:1–14
- Matsumoto Y, Hiramatsu C, Matsushita Y et al (2014) Evolutionary renovation of L/M opsin polymorphism confers a fruit discrimination advantage to ateline New World monkeys. Mol Ecol 7:1799–1812
- Melin AD, Fedigan LM, Hiramatsu C, Sendall CL, Kawamura S (2007) Effects of colour vision phenotype on insect capture by a freeranging population of white-faced capuchins, *Cebus capucinus*. Anim Behav 73:205–214
- Melin AD, Fedigan LM, Hiramatsu C, Kawamura S (2008) Polymorphic color vision in white-faced capuchins (*Cebus capucinus*): is there foraging niche divergence among phenotypes? Behav Ecol Sociobiol 62:659–670
- Melin AD, Fedigan LM, Young HC, Kawamura S (2010) Can color vision variation explain sex differences in invertebrate foraging by capuchin monkeys? Curr Zool 56:300–312
- Melin AD, Moritz GL, Fosbury RAE, Kawamura S, Dominy NJ (2012) Why aye-ayes see blue. Am J Primatol 74:185–192
- Melin AD, Matsushita Y, Moritz G, Dominy NJ, Kawamura S (2013) Inferred M/L cone opsin polymorphism of ancestral tarsiers sheds dim light on the origin of anthropoid primates. Proc R Soc B 208: 1759
- Melin AD, Chiou K, Walco E, Bergstrom M, Kawamura S, Fedigan LM (2017a) Trichromacy increases fruit intake rates of wild capuchins (*Cebus capucinus imitator*). Proc Natl Acad Sci U S A 114:10402– 10407
- Melin AD, Khetpal V, Matsushita Y, Zhou K, Campos FA, Welker B, Kawamura S (2017b) Howler monkey foraging ecology suggests convergent evolution of routine trichromacy as an adaptation for folivory. Ecol Evol 7:1421–1434
- Mittermeier RA, Louis EE, Richardson M et al (2010) Lemurs of Madagascar, 3rd edn. Conservation International, Arlington
- Nathans J (1999) The evolution and physiology of human color vision: insights from molecular genetic studies of visual pigments. Neuron 24:299–312
- Neitz M, Neitz J, Jacobs GH (1991) Spectral tuning of pigments underlying red-green color vision. Science 252:971–974

- Nevo O, Valenta K, Razafimandimby D, Melin AD, Ayasse M, Chapman CA (2018) Frugivores and the evolution of fruit colour. Biol Lett 14: 20180377
- Nsubuga AM, Robbins MM, Roeder AD, Morin PA, Boesch C, Vigilant L (2004) Factors affecting the amount of genomic DNA extracted from ape faeces and the identification of an improved sample storage method. Mol Ecol 13:2089–2094
- Olsson P, Lind O, Kelber K (2018) Chromatic and achromatic vision: parameter choice and limitations for reliable model predictions. Behav Ecol 29:273–282
- Osorio D, Vorobyev M (2018) Principles and application of the receptor noise model of color discrimination: a comment on Olsson et al. Behav Ecol 29:283–284
- Osorio D, Ruderman DL, Cronin TW (1998) Estimation of errors in luminance signals encoded by primate retina resulting from sampling of natural images with red and green cones. J Opt Soc Am A 15:16–22
- Osorio D, Smith AC, Vorobyev M, Buchanan-Smith HM (2004) Detection of fruit and the selection of primate visual pigments for color vision. Am Nat 164:696–708
- Overdorff DJ (1996) Ecological correlates to activity and habitat use of two prosimian primates: *Eulemur rubriventer* and *Eulemur fulvus rufus* in Madagascar. Am J Primatol 40:327–342
- Pariente GF (1980) Quantitative and qualitative study of the light available in the natural biotope of Malagasy prosimians. In: Charles-Dominique P, Cooper HM, Hladik A, Hladik CM, Pages E, Pariente GF, Petter-Rousseaux A, Petter JJ, Schilling A (eds) Nocturnal Malagasy primates: ecology, physiology and behaviour. Academic Press, New York, pp 117–134
- Pessoa D, Maia R, Ajuz R, Moraes P, Spyrides M, Pessoa V (2014) The adaptive value of primate color vision for predator detection. Am J Primatol 76:721–729
- R Core Team (2014) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna https:// www.R-project.org/
- R Core Team (2017) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna https:// www.R-project.org/
- Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. Conserv Biol 17:230–237
- Regan BC, Julliot C, Simmen B, Vienot F, Charles-Dominique P, Mollon JD (2001) Fruits, foliage and the evolution of primate colour vision. Philos Trans R Soc B 356:229–283
- Rennison DJ, Owens GL, Taylor GS (2012) Opsin gene duplication and divergence in ray-finned fish. Mol Phylogenet Evol 62:986–1008
- Revell LJ (2012) phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol Evol 3:217–223
- Roth LSV, Balkenius A, Kelber A (2008) The absolute threshold of colour vision in the horse. PLoS One 3:e3711

- Sambrook J, Fritch EF, Maniatus T (1989) Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor Press, New York
- Smith AC, Surridge AK, Prescott MJ, Osorio D, Mundy NI, Buchanan-Smith HM (2012) The effect of colour vision status on insect prey capture efficiency by captive and wild tamarins (*Saguinus* spp.). Anim Behav 83:479–486
- Springer MS, Meredith RW, Gatesy J, Emerling CA, Park J, Rabosky DL, Stadler T, Steiner C, Ryder OA, Janečka JE, Fisher CA, Murphy WJ (2012) Macroevolutionary dynamics and historical biogeography of primate diversification inferred from a species supermatrix. PLoS One 7:e49521
- Sumner P, Mollon JD (2000) Catarrhine photopigments are optimized for detecting targets against a foliage background. J Exp Biol 203: 1963–1986
- Surridge AK, Osorio D, Mundy NI (2003) Evolution and selection of trichromatic vision in primates. Trends Ecol Evol 18:198–205
- Tan Y, Li WH (1999) Vision trichromatic vision in prosimians. Nature 402(36):36
- Valenta K, Burke RJ, Styler SA, Jackson DA, Melin AD, Lehman SM (2013) Colour and odour drive fruit selection and seed dispersal by mouse lemurs. Sci Rep 3:2424
- Valenta K, Edwards M, Rafaliarison RR, Johnson SE, Holmes SM, Brown KA, Dominy NJ, Lehman SM, Parra EJ, Melin AD (2016) Visual ecology of true lemurs suggests a cathemeral origin for the primate cone opsin polymorphism. Funct Ecol 30:932–942
- Veilleux CC, Bolnick DA (2009) Opsin gene polymorphism predicts trichromacy in a cathemeral lemur. Am J Primatol 71:86–90
- Veilleux CC, Cummings ME (2012) Nocturnal light environments and species ecology: implications for nocturnal color vision in forests. J Exp Biol 215:4085–4096
- Veilleux CC, Jacobs RL, Cummings ME, Louis EE, Bolnick DA (2014) Opsin genes and visual ecology in a nocturnal folivorous lemur. Int J Primatol 35:88–107
- Veilleux CC, Scarry CJ, Di Fiore A, Kirk EC, Bolnick DA, Lewis RJ (2016) Group benefit associated with polymorphic trichromacy in a Malagasy primate (*Propithecus verreauxi*). Sci Rep 6:38418
- Vogel ER, Neitz M, Dominy NJ (2007) Effect of color vision phenotype on the foraging of wild white-faced capuchins, *Cebus capucinus*. Behav Ecol 18:292–297
- Williams AJ, Hunt DM, Bowmaker JK, Mollon JD (1992) The polymorphic photopigments of the marmoset: spectral tuning and genetic basis. EMBO J 11:2039–2045
- Wright PC (1992) Primate ecology, rainforest conservation, and economic development: building a national park in Madagascar. Evol Anthropol 1:25–33
- Wyszecki G, Stiles WS (1982) Color science: concepts and methods, quantitative data and formulae, 2nd edn. Wiley, New York
- Yamashita N, Stoner K, Riba-Hernández P, Dominy N, Lucas P (2005) Light levels used during feeding by primate species with different color vision phenotypes. Behav Ecol Sociobiol 58:618–629